



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS FOR SOLUTION.

ALGEBRA.

306. Proposed by J. C. CORBIN, Pine Bluff, Ark.

Muir gives the following problem:

$$\text{Prove: } \begin{vmatrix} 1 & a & a & a^2 \\ 1 & b & b & b^2 \\ 1 & c & c' & cc' \\ 1 & d & d' & dd' \end{vmatrix} = (a-b) \begin{vmatrix} 1 & ab & a+b \\ 1 & cd' & c+d' \\ 1 & c'd & c'+d \end{vmatrix}$$

which, of course, can be solved by finding the terms of both determinants. Is there any method of changing from one form to the other which is direct?

307. Proposed by J. SCHEFFER, Hagerstown, Md.

If $y^x=2$ and $x^y=3$, find x and y .

GEOMETRY.

339. Proposed by G. E. BROCKWAY, Boston, Mass.

Of all triangles that can be inscribed in a given triangle, that formed by joining the feet of the altitudes has the minimum perimeter. Prove by means of the straight line and circle.

340. Proposed by J. H. MEYERS, S. J., Sacred Heart College, Augusta, Ga.

Given trapezoid $ABCD$. Prolong AB and CD , the non-parallel sides, to meet in E . On AE as diameter construct semi-circle $ALGE$. With BE as radius construct arc BG . Draw GH perpendicular to AE . Bisect AH at K . Erect KL perpendicular to AE . Construct arc LM with LE as radius. Draw MN perpendicular to DC . Prove that MN bisects the trapezoid $ABCD$, angles ADC and BCD being right angles.

CALCULUS.

265. Proposed by V. M. SPUNAR, M. and E. E., East Pittsburg, Pa.

Find two curves which possess the property that the tangents TP and TQ to the inner one always make equal angles with the tangent TT' to the outer.

266. Proposed by C. N. SCHMALL, New York City.

Show that the n th derivative of the fraction u/v can be expressed in the form of a determinant, u and v being functions of x .

MECHANICS.

221. Proposed by W. J. GREENSTREET, Stroud, England.

Two smooth intersecting planes are each at 45° to the horizon. Between them lies a cylinder of elliptic cross section. Find the position of equilibrium.